

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

**0 380 257**  
**A2**

(12)

# EUROPEAN PATENT APPLICATION

(21) Application number: 90300603.9

(22) Date of filing: 22.01.90

(51) Int. Cl.<sup>5</sup>: **C07D 487/04, C07D 471/04,**  
**A61K 31/50, C07D 237/28,**  
**/(C07D487/04,237:00,209:00),**  
**(C07D471/04,237:00,221:00)**

The title of the invention has been amended  
(Guidelines for Examination in the EPO, A-III,  
7.3).

(30) Priority: 23.01.89 GB 8901430

(43) Date of publication of application:  
01.08.90 Bulletin 90/31

(84) Designated Contracting States:  
CH DE FR GB IT LI NL


(71) Applicant: **SMITHKLINE BECKMAN**  
**INTERCREDIT B.V.**  
28-34 Blaak P.O. Box 2  
NL-3000 DG Rotterdam(NL)

(72) Inventor: **Brown, Thomas Henry**  
**Smith Kline & French Research Ltd., The**  
**Frythe**  
**Welwyn, Hertfordshire AL6 9AR(GB)**  
Inventor: **Blurton, Peter**  
**Smith Kline & French Research Ltd., The**  
**Frythe**  
**Welwyn, Hertfordshire AL6 9AR(GB)**

(74) Representative: **Giddings, Peter John, Dr. et al**  
**SmithKline Beecham, Corporate Patents,**  
**Mundells**  
**Welwyn Garden City, Hertfordshire AL7**  
**1EY(GB)**

(54) Condensed cinnoline derivatives, process and intermediates for their preparation and pharmaceutical compositions containing them.


(57) Compounds of structure (I)



(I)

in which R<sup>1</sup> and R<sup>2</sup> are the same or different and are each hydrogen, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy, phenyl, C<sub>1-6</sub>alkylthio, C<sub>1-6</sub>alkanoyl, amino, C<sub>1-6</sub>alkylamino, diC<sub>1-6</sub>alkylamino, halogen or trifluoromethyl;  
R<sup>3</sup> is hydrogen, C<sub>1-6</sub>alkyl or a group

EP 0 380 257 A2




R<sup>4</sup> to R<sup>6</sup> are the same or different and are each hydrogen, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy, C<sub>1-6</sub>alkylthio, halogen, cyano, amino, hydroxy, carbamoyl, carboxy, C<sub>1-6</sub>alkanoyl or trifluoromethyl; n is 0 to 4; and A is  $-(CH_2)_2-$ ,  $-(CH_2)_3-$  or  $-CH=CH-$ ; processes for their preparation, pharmaceutical compositions containing them and their use in therapy as inhibitors of gastric acid secretion.

## Compounds


The present invention relates to substituted cinnoline derivatives, processes for their preparation, intermediates useful in their preparation, pharmaceutical compositions containing them and their use in therapy.

Accordingly the present invention provides, in a first aspect compounds of structure (I)



in which

R<sup>1</sup> and R<sup>2</sup> are the same or different and are each hydrogen, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy, phenyl, C<sub>1-6</sub>alkylthio, C<sub>1-6</sub>alkanoyl, amino, C<sub>1-6</sub>alkylamino, diC<sub>1-6</sub>alkylamino, halogen or trifluoromethyl;  
 R<sup>3</sup> is hydrogen, C<sub>1-6</sub>alkyl or a group




R<sup>4</sup> to R<sup>6</sup> are the same or different and are each hydrogen, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy, C<sub>1-6</sub>alkylthio, halogen, cyano, amino, hydroxy, carbamoyl, carboxy, C<sub>1-6</sub>alkanoyl or trifluoromethyl;  
 n is 0 to 4; and


A is -(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-, or -CH=CH-;  
 or a salt thereof.

Suitably, R<sup>1</sup> and R<sup>2</sup> are the same or different and are each, hydrogen, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy, phenyl, C<sub>1-6</sub>alkylthio, C<sub>1-6</sub>alkanoyl, amino, C<sub>1-6</sub>alkylamino, diC<sub>1-6</sub>alkylamino, halogen or trifluoromethyl. Preferably, one of R<sup>1</sup> and R<sup>2</sup> is hydrogen and the other is hydrogen or C<sub>1-6</sub>alkyl.

Suitably, R<sup>3</sup> is hydrogen, C<sub>1-6</sub>alkyl or



Preferably, R<sup>3</sup> is C<sub>1-6</sub>alkyl or



Suitably R<sup>4</sup> to R<sup>6</sup> are the same or different and are each hydrogen, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy, C<sub>1-6</sub>alkylthio, halogen, cyano, amino, hydroxy, carbamoyl, carboxy, C<sub>1-6</sub>alkanoyl or trifluoromethyl. Preferably one of R<sup>4</sup> to R<sup>6</sup> is hydrogen and the other two are the same or different and are hydrogen, C<sub>1-6</sub>alkyl or hydroxy.


Suitably n is 0 to 4; preferably n is 0.

Suitably A is  $-(CH_2)_3-$ ,  $-(CH_2)_2$  or  $CH=CH$ ; preferably A is  $-(CH_2)_2-$ .

C<sub>1-6</sub>alkyl groups (either alone or as part of another group) can be straight or branched.

It will be appreciated that compounds of structure (I) in which one or more of R<sup>1</sup> to R<sup>6</sup> is a C<sub>3-6</sub>alkyl group (either alone or as part of another group) may contain an asymmetric centre due to the presence of the C<sub>3-6</sub>alkyl group. Such compounds will exist as optical isomers (enantiomers). Both the pure enantiomers, racemic mixtures (50% of each enantiomer) and unequal mixtures of the two are included within the scope of the present invention. Further, all diastereomeric forms possible (pure enantiomers and mixtures thereof) are within the scope of the invention.

The present invention provides in a further aspect a process for the preparation of a compound of structure (I) or a pharmaceutically acceptable salt thereof which comprises cyclisation of a compound of structure (II)



in which R<sup>1</sup> and R<sup>2</sup> are as described for structure (I); m is 2 or 3 and X<sup>1</sup> and X<sup>2</sup> are leaving groups, in the presence of a compound of structure R<sup>3</sup>NH<sub>2</sub> or R<sup>3</sup>NH<sub>3</sub><sup>+</sup>Cl<sup>-</sup> (III) in which R<sup>3</sup> is as described for structure (I); or

- oxidising a compound in which A is  $(CH_2)_2$  to a compound in which A is  $-CH=CH-$ ;
- forming a salt.

Suitable leaving groups X<sup>1</sup> and X<sup>2</sup> will be apparent to those skilled in the art and include for example halogen, in particular chlorine.


Suitably the cyclisation of a compound of structure (II) in the presence of a compound of structure (III) is carried out in a solvent and at a temperature of between ambient and the reflux temperature of the solvent used. Preferably the reaction is carried out at elevated temperature in a suitable solvent. For example, in a solvent such as 1,4-Dioxan or phenol at elevated temperature. Suitably the compound (III) is present as the free base, preferably in the protonated form R<sup>3</sup>NH<sub>3</sub><sup>+</sup>Cl<sup>-</sup>.

Suitable reagents for carrying out the oxidation of compounds of structure (I) in which A is  $(CH_2)_2$  to compounds of structure (I) in which A is  $CH=CH$  will be apparent to those skilled in the art and include, for example, dehydrogenation in the presence of a noble metal catalyst, in particular palladium-on-carbon. Under certain conditions, as described in the examples hereinafter, the dehydrogenation can also take place in the absence of a solid catalyst.

Acid addition salts of the compounds of structure (I) in particular pharmaceutically acceptable salts can be prepared by standard procedures by, for example, reaction with suitable organic and inorganic acids the nature of which will be apparent to persons skilled in the art. For example, pharmaceutically acceptable salts can be formed by reaction with hydrochloric, sulphuric, or phosphoric acids; aliphatic, aromatic or heterocyclic sulphonic acids or carboxylic acids such as, for example, citric, maleic or fumaric acids.

The intermediate compounds of structure (II) are new and as such form a further aspect of the present invention. They can be prepared by procedures analogous to those known in the art. For example, compounds of structure (II) in which both X<sup>1</sup> and X<sup>2</sup> are chloro can be prepared by the reaction sequence outlined in Scheme 1.

Scheme I



It is to be noted, and apparent to those skilled in the art that in the foregoing reactions, where necessary, groups R<sup>1</sup> to R<sup>6</sup> will be in protected form when undergoing the procedures described.

The compounds of structure (I) and their pharmaceutically acceptable salts and related compounds for which no pharmaceutical utility has been previously disclosed exert an anti-secretory effect by inhibition of the gastrointestinal  $\text{H}^+\text{K}^+\text{ATPase}$  enzyme (Fellenius E., Berglindh T., Sachs G., Olke L., Elander B., Sjostrand S.E., and Wahlmark B., 1981, *Nature*, 290, 159-61).

In a further aspect therefore the present invention provides the use of the compounds of structure (I) and pharmaceutically acceptable salts thereof in therapy. The compounds of structure (I) and their pharmaceutically acceptable salts inhibit exogenously and endogenously stimulated gastric acid secretion and are useful in the treatment of gastrointestinal diseases in mammals, in particular humans. Such diseases include, for example, gastric and duodenal ulcers, and Zollinger-Ellison Syndrome.

Further, the compounds of structure (I) can be used in the treatment of other disorders where an anti-secretory effect is desirable for example in patients with gastritis, NSAID induced gastritis, gastric ulcers, acute upper intestinal bleeding, in patients with a history of chronic and excessive alcohol consumption, and in patients with gastro oesophageal reflux disease (GERD).

In therapeutic use, the compounds of the present invention are usually administered in a standard pharmaceutical composition. The present invention therefore provides, in a further aspect pharmaceutical compositions comprising a compound of structure (I) or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

The compounds of structure (I) and their pharmaceutically acceptable salts which are active when given orally can be formulated as liquids, for example syrups, suspensions or emulsions, tablets, capsules and lozenges.

A liquid formulation will generally consist of a suspension or solution of the compound or pharmaceutically acceptable salt in a suitable liquid carrier(s) for example, ethanol, glycerine, non-aqueous solvent, for example polyethylene glycol, oils, or water with a suspending agent, preservative, flavouring or colouring agent.

A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s)

routinely used for preparing solid formulations. Examples of such carriers include magnesium stearate, starch, lactose, sucrose and cellulose.

A composition in the form of a capsule can be prepared using routine encapsulation procedures. For example, pellets containing the active ingredient can be prepared using standard carriers and then filled into a hard gelatin capsule; alternatively, a dispersion or suspension can be prepared using any suitable pharmaceutical carrier(s), for example aqueous gums, celluloses, silicates or oils and the dispersion or suspension then filled into a soft gelatin capsule.

Typical parenteral compositions consist of a solution or suspension of the compound or pharmaceutically acceptable salt in a sterile aqueous carrier or parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis oil or sesame oil. Alternatively, the solution can be lyophilised and then reconstituted with a suitable solvent just prior to administration.

A typical suppository formulation comprises a compound of structure (I) or a pharmaceutically acceptable salt thereof which is active when administered in this way, with a binding and/or lubricating agent such as polymeric glycols, gelatins or cocoa butter or other low melting vegetable or synthetic waxes or fats.

Preferably the composition is in unit dose form such as a tablet or capsule.

Each dosage unit for oral administration contains preferably from 1 to 250 mg (and for parenteral administration contains preferably from 0.1 to 25 mg) of a compound of structure (I) or a pharmaceutically acceptable salt thereof calculated as the free base.

The pharmaceutically acceptable compounds of the invention will normally be administered to a subject for the treatment of gastro-intestinal diseases and other conditions caused or exacerbated by gastric acidity. The daily dosage regimen for an adult patient may be, for example, an oral dose of between 1 mg and 500 mg, preferably between 1 mg and 250 mg, or an intravenous, subcutaneous, or intramuscular dose of between 0.1 mg and 100 mg, preferably between 0.1 mg and 25 mg; of the compound of structure (I) or a pharmaceutically acceptable salt thereof calculated as the free base, the compound being administered 1 to 4 times per day. Suitably the compounds will be administered for a period of continuous therapy, for example for a week or more.

In addition, the compounds of the present invention can be co-administered with further active ingredients, such as antacids (for example magnesium carbonate or hydroxide and aluminium hydroxide), non-steroidal anti-inflammatory drugs (for example indomethacin, aspirin or naproxen), steroids, or nitrite scavengers (for example ascorbic acid or aminosulphonic acid), or other drugs used for treating gastric ulcers (for example pirenzepine, prostanoids for example 16,16 dimethyl PGE<sub>2</sub>, or histamine H<sub>2</sub>-antagonists (for example, cimetidine).

The following examples illustrate the invention. Temperatures are recorded in degrees centigrade.

#### Example 1

##### 1-(2-Methylphenyl)-2,3-dihydropyrrolo[3,2-c]cinnoline

##### a) 1-(2-Aminophenyl)-4-chlorobutan-1-one hydrochloride

To a solution of boron trichloride (55 g) in 1,2-dichloroethane (200 ml) at 0° was added dropwise a solution of aniline (39.1 g, 0.42 mol) in 1,2-dichloroethane (50 ml) maintaining the temperature below 5°. On completion of the addition 4-chlorobutyronitrile (41.4 g, 0.42 mol) and aluminium chloride (53.34 g, 0.42 mol) were added successively and the reaction mixture allowed to warm up to room temperature then heated under reflux for 20 hours. After allowing to cool, 2N hydrochloric acid (100 ml) was added and the reaction mixture heated under reflux for 0.5 hour. The solid obtained was collected by filtration, taken up in water and extracted with chloroform (5 x 200 ml). The chloroform extracts were combined, washed with dilute base, then water, dried over magnesium sulphate, filtered and evaporated under reduced pressure to give an oil. The oil was dissolved in ether and ethanolic HCl was added, the resulting solid was collected by filtration and dried, 28.5 g. Recrystallization from ethanol/diethyl ether gave the product, m.p. 152-154°

C <sub>10</sub> H <sub>12</sub> ClNO HCl	
Found	C 51.27, H 5.52, N 6.02, Cl 15.11, Cl <sup>-</sup> 15.10%
Requires	C 51.30, H 5.60, N 5.98, Cl 15.15, Cl <sup>-</sup> 15.15%

5

10 b) 3-(2-Chloroethyl)-4-oxo-(1H)-cinnoline.

A solution of 1-(2-aminophenyl)-4-chlorobutan-1-one hydrochloride (11.3 g, 0.0427 mol) in concentrated hydrochloric acid (1200 ml) was cooled down to 0° with stirring. Then sodium nitrite (3.54 g, 0.051 mol) in water (25 ml) was added dropwise. On completion of the addition, the reaction mixture was allowed to warm to room temperature and stirred overnight. The solution was evaporated under reduced pressure to dryness. The solid obtained was washed with a solution of sodium acetate and collected by filtration to give the title compound, 5.32 g. Recrystallization from methanol gave the product, m.p. 201-203°.

20

C <sub>10</sub> H <sub>9</sub> ClN <sub>2</sub> O	
Found	C 57.53, H 4.23, N 13.28, Cl 16.84%
Requires	C 57.56, H 4.35, N 13.43, Cl 16.99%

25

c) 3-(2-Chloroethyl)-4-chlorocinnoline.

30 3-(2-Chloroethyl)-4-oxo-(1H)-cinnoline (9 g, 0.043 mol) and phosphorus oxychloride (90 ml) were heated under reflux for 1 hour. After allowing to cool, the excess phosphorus oxychloride was evaporated under reduced pressure and the residue was poured onto ice (100 g), basified with concentrated ammonia solution, and extracted with chloroform (3 x 100 ml). The chloroform extracts were combined, dried over magnesium sulphate and evaporated under reduced pressure to give a solid, 8.00 g. Recrystallization from diethyl ether gave the title compound, m.p. 81-82°.

35

C <sub>10</sub> H <sub>8</sub> Cl <sub>2</sub> N <sub>2</sub>	
Found	C 52.98, H 3.68, N 12.21, Cl 30.82%
Requires	C 52.89, H 3.55, N 12.34, Cl 31.23%

40

45 d) 1-(2-Methylphenyl)-2,3-dihydropyrrolo[3,2-c]cinnoline.

4-Chloro-3-(2-chloroethyl)cinnoline (1.52 g, 0.0067 mol) and o-toluidine (1.43 g, 0.0134 mol) in 1,4 dioxan (20 ml) were heated under reflux for 20 hours. The solvent was evaporated under reduced pressure and the residue was partitioned between chloroform and dilute sodium bicarbonate solution. The aqueous layer was extracted with chloroform (6 x 50 ml). The chloroform extracts were combined, dried over magnesium sulphate, filtered and evaporated under reduced pressure to give a brown oil. The oil was purified by chromatography using chloroform (1 litre), chloroform/methanol (20:1) as eluant. The fractions containing the product were combined and evaporated under reduced pressure to give an oil. On trituration with diethyl ether a solid was obtained. The solid was collected by filtration and dried, 0.76 g. Recrystallization from ethyl acetate gave the title compound, 0.48 g, m.p. 140-142°.

55

C <sub>17</sub> H <sub>15</sub> N <sub>3</sub>	
Found	C 77.71, H 5.81, N 16.01%
Requires	C 78.13, H 5.78, N 16.08%

### Example 2

#### 1-(2-methylphenyl)-6-methyl-2,3-dihydropyrrolo[3,2-c]cinnoline

##### a) 1-(2-Amino-3-methylphenyl)-4-chlorobutan-1-one hydrochloride.

To a solution of boron trichloride (55 g) in 1,2 dichloroethane at 0° was added dropwise a solution of o-toluidine (45.7 g, 0.426 mol) in 1,2 dichloroethane (50 ml) maintaining the temperature below 5°. On completion of the addition 4-chlorobutyronitrile (41.4 g, 0.426 mol) and aluminium chloride (53.34 g, 0.426 mol) were added successively. The reaction mixture was allowed to warm up to room temperature and stirred for 1.5 hours, then heated under reflux for 20 hours. The reaction mixture was then cooled to room temperature poured onto ice (500 g) and 2N hydrochloric acid (200 ml). The mixture was then heated under reflux for 0.5 hour. The organic layer was separated and the aqueous layer was extracted with chloroform (3 x 100 ml). The organic extracts were combined, washed with water, dried over magnesium sulphate, filtered and evaporated under reduced pressure to give an oil. The oil was dissolved in diethyl ether and HCl gas bubbled through. The resulting solid was collected by filtration and dried (57 g). Recrystallization from ethanol/diethyl ether gave the title compound, m.p. 148-150°.

C <sub>11</sub> H <sub>14</sub> ClNO HCl					
Found	C 53.25,	H 6.03,	N 5.64,	Cl 14.09,	Cl <sup>-</sup> 14.19%
Requires	C 53.24	H 6.09,	N 5.64,	Cl 14.19,	Cl <sup>-</sup> 14.19%

##### b) 3-(2-Chloroethyl)-8-methyl-4-oxo-(1H)-cinnoline.

To a solution of 1-(2-amino-3-methylphenyl)-4-chlorobutan-1-one hydrochloride (20 g, 0.081 Mol) and concentrated hydrochloric acid (1000 ml) at 0° was added dropwise a solution of sodium nitrite (6.07 g, 0.088 mol) in water (50 ml) maintaining the temperature below 5°. The reaction mixture was stirred at room temperature for 20 hours. The solvent was evaporated under reduced pressure to give a solid. The solid was washed with sodium acetate solution and collected by filtration and dried, 13.5 g. Recrystallization from methanol gave the title compound, m.p. 222-225°.

C <sub>11</sub> H <sub>11</sub> ClN <sub>2</sub> O	
Found	C 59.27, H 4.98, N 12.50, Cl 15.04%
Requires	C 59.33, H 4.98, N 12.58, Cl 14.98%

##### c) 3-(2-Chloroethyl)-4-chloro-8-methylcinnoline.

3-(2-Chloroethyl)-8-methyl-4-oxo-(1H)-cinnoline (9 g, 0.04 mol) and phosphorus oxychloride (100 ml)



were heated under reflux for 2 hours. The excess phosphorus oxychloride was evaporated under reduced pressure and the residue was poured onto ice, basified with concentrated ammonia solution and extracted with chloroform (3 x 200 ml). The chloroform extracts were combined, dried over magnesium sulphate, filtered and evaporated under reduced pressure to give a dark brown solid. Recrystallization from diethyl ether/petroleum ether (b.p. 40-60°) gave an analytically pure sample, m.p. 102-104°.

C <sub>11</sub> H <sub>10</sub> Cl <sub>2</sub> N <sub>2</sub>	
Found	C 55.12, H 4.09, N 11.78, Cl 29.41%
Requires	C 54.79, H 4.18, N 11.62, Cl 29.41%

d) 1-(2-Methylphenyl)-6-methyl-2,3-dihydropyrrolo[3,2-c]cinnoline.

3-(2-Chloroethyl)-4-chloro-8-methylcinnoline (2.5 g, 0.01 mol) and o-toluidine hydrochloride (1.4 g, 0.01 mol) in 1,4 dioxan (20 ml) were heated under reflux for 20 hours. The solvent was evaporated under reduced pressure to give an oil. The oil was partitioned between chloroform and dilute sodium carbonate solution and extracted further with chloroform (4 x 100 ml). The chloroform extracts were combined, dried over magnesium sulphate, filtered and evaporated to give an oil. The oil was purified by column chromatography using dichloromethane then chloroform as eluant. The solid obtained was recrystallized from diethyl ether/ dichloromethane to give the title compound, 0.6 g, m.p. 135-137°.

C <sub>18</sub> H <sub>17</sub> N <sub>3</sub>	
Found	C 78.49, H 6.31, N 15.30%
Requires	C 78.52, H 6.22, N 15.26%

Example 3

1-(2-Methylphenyl)-6-methylpyrrolo[3,2-c]cinnoline.

To a refluxing slurry of 10%Pd/C (0.4 g) in diphenyl ether (20 ml) was added 1-(2-methylphenyl)-6-methyl-2,3-dihydropyrrolo[3,2-c] cinnoline (1.25 g, 0.0045 mol). After 0.5 hour at reflux temperature the reaction mixture was cooled, diluted with chloroform and the catalyst was filtered off over hyflo. The product was isolated by column chromatography using dichloromethane (1l) then chloroform. The oil obtained was crystallized from ethyl acetate, 0.42 g, m.p. 105-106°.

C <sub>18</sub> H <sub>15</sub> N <sub>3</sub>	
Found	C 79.10, H 5.61, N 15.48%
Requires	C 79.09, H 5.53, N 15.37%

Example 4

1-(4-Hydroxy-2-methylphenyl)-2,3-dihydropyrrolo[3,2-c]cinnoline hydrochloride

4-Chloro-3-(2-chloroethyl)cinnoline (1 g, 0.0044 mol) and 4-hydroxy-2-methylphenylamine (0.54 g, 0.0044 mol) in 1,4 dioxan (20 ml) were heated under reflux for 20 hours. The solid obtained was collected by filtration and dried. Recrystallization from ethanol gave the title compound, 0.4 g, m.p.  $>300^{\circ}$

C <sub>17</sub> H <sub>15</sub> N <sub>3</sub> O HCl	
Found	C 64.67, H 5.18, N 13.14, Cl <sup>-</sup> 11.16%
Requires	C 65.07, H 5.14, N 13.39, Cl <sup>-</sup> 11.30%

#### Example 5

##### N-propyl-2,3-dihydropyrrolo[3,2-c]cinnoline

3-(2-Chloroethyl)-4-chlorocinnoline (2 g, 0.088 mol), propylamine (4 g, 0.067 mol) and phenol (2 g, 0.0212 mol) were heated in an oil bath at  $160^{\circ}$  for 20 hours. The reaction mixture was diluted with chloroform (200 ml) and extracted with 2N sodium hydroxide (3 x 100 ml). The chloroform extracts were then washed with water (50 ml), dried over magnesium sulphate, filtered and evaporated under reduced pressure to give an oil. The oil was purified by chromatography using a gradient elution chloroform/methanol (0-5% MeOH). The fractions containing the product were combined and evaporated to give a yellow solid, 1.4 g. Recrystallization from ethyl acetate gave the title compound, 1.21 g, m.p.  $116-118^{\circ}$ .

C <sub>13</sub> H <sub>15</sub> N <sub>3</sub>	
Found	C 73.20, H 7.10, N 19.72%
Requires	C 73.21, H 7.09, N 19.70%

#### Example 6

##### N-Benzylpyrrolo[3,2-c]cinnoline

3-(2-Chloroethyl)-4-chlorocinnoline (2 g, 0.0088 mol), benzylamine (4 g, 0.037 mol) and phenol (2 g, 0.0212 mol) were heated in an oil bath at  $160^{\circ}$  for 20 hours. The reaction had not gone to completion so the temperature was raised to  $190^{\circ}$  for a further 6 hours. The reaction mixture was dissolved in chloroform (300 ml) and extracted with 2N sodium hydroxide (3 x 100 ml). The chloroform extracts were washed with water (100 ml), dried over magnesium sulphate, filtered and evaporated under reduced pressure to give an oil. The oil was purified by chromatography using chloroform as eluant. The fractions containing the product were combined and evaporated to dryness to give a brown solid. Recrystallization from isopropyl alcohol gave the title compound, 0.5 g, m.p.  $190-192^{\circ}$ .

C <sub>17</sub> H <sub>13</sub> N <sub>3</sub>	
Found	C 78.72, H 5.02, N 16.01%
Requires	C 78.74, H 5.05, N 16.21%

Example 7

5 1-(4-Hydroxy-2-methylphenyl)-6-methyl-2,3-dihydropyrrolo[3,2-c]cinnoline hydrochloride.

3-(2-Chloroethyl)-4-chloro-8-methylcinnoline (2 g, 0.0083 mol) and 4-hydroxy-2-methylaniline hydrochloride (1.32 g, 0.0083 mol) in isopropanol (50 ml) were heated under reflux in a nitrogen atmosphere for 20 hours. The solid obtained was collected by filtration and dried. Recrystallization from isopropanol/ethanolic  
 10 HCl gave the title compound as a partial hydrate, 0.64 g, m.p. 292-294°.

C <sub>18</sub> H <sub>17</sub> N <sub>3</sub> O HCl 0.6H <sub>2</sub> O	
Found	C 63.69, H 5.73, N 12.39, Cl <sup>-</sup> 10.63%
Requires	C 63.84, H 5.71, N 12.40, Cl <sup>-</sup> 10.47%

20

Example A

A tablet for oral administration is prepared by combining

25

	Mg/Tablet
Compound of structure (I)	100
Mannitol	153
Starch	33
crospovidone	12
microcrystalline cellulose	30
magnesium stearate	2
	<u>330</u>
	mg

35

into a 9 mm tablet.

40

Example B

An injection for parenteral administration was prepared from the following

45

Compound of Structure 1	6.68%(w:v)
1M citric acid	30%(v:v)
sodium hydroxide (qs)	to pH 3.2
water for injection EP	to 100 ml

50

The compound of Structure 1 was dissolved in the citric acid and the pH slowly adjusted to pH 3.2 with the sodium hydroxide solution. The solution was then made up to 100 ml with water, sterilised by filtration  
 55 and sealed into appropriately sized ampoules and vials.

Biological Data.

(A) H<sup>+</sup>K<sup>+</sup>ATPase Activity.

The effects of a single high concentration (100  $\mu$ M) of a compound of structure (I) on K<sup>+</sup>-stimulated ATPase activity in lyophilised gastric vesicles was determined. Preferred compounds of structure (I) were also tested over a range of concentrations to determine IC<sub>50</sub> values.

(i) Preparation of lyophilised gastric vesicles (H/K-ATPase).

Lyophilised gastric vesicles were prepared from pig fundic mucosa after the method of Keeling et. al. (Biochem. Pharmacol., 34, 2967, 1985).

(ii) K<sup>+</sup>-stimulated ATPase activity.

K<sup>+</sup>-stimulated ATPase activity was determined at 37° in the presence of the following : 10 mM Pipes/Tris buffer pH 7.0, 2 mM MgSO<sub>4</sub>, 1 mM KCl, 2 mM Na<sub>2</sub>ATP and 3  $\mu$ g protein/ml lyophilised gastric vesicles. After incubation for 30 minutes, the inorganic phosphate hydrolysed from ATP was determined by the method of Yoda and Hokin (Biochem. Biophys. Res. Commun. 40, 880, 1970).

Compounds of structure (I) were dissolved in dimethylsulphoxide which up to the highest concentration used had no effect on K<sup>+</sup>-stimulated ATPase activity.


The effect of the highest concentration of each compound of structure (I) on the recovery of a standard amount of inorganic phosphate was also determined.

(iii) Results.

The compounds of each of examples 1 to 5 were assayed in the above-noted screen and found to have IC<sub>50</sub> values in the range of from 1.8 to 11.1  $\mu$ M.

**Claims**


1. A compound of structure (I)



in which

R<sup>1</sup> and R<sup>2</sup> are the same or different and are each hydrogen, C<sub>1</sub>-<sub>6</sub>alkyl, C<sub>1</sub>-<sub>6</sub>alkoxy, phenyl, C<sub>1</sub>-<sub>6</sub>alkylthio, C<sub>1</sub>-<sub>6</sub>alkanoyl, amino, C<sub>1</sub>-<sub>6</sub>alkylamino, diC<sub>1</sub>-<sub>6</sub>alkylamino, halogen or trifluoromethyl;

R<sup>3</sup> is hydrogen, C<sub>1</sub>-<sub>6</sub>alkyl or a group



10  $R^4$  to  $R^6$  are the same or different and are each hydrogen,  $C_1-6$  alkyl,  $C_1-6$  alkoxy,  $C_1-6$  alkylthio, halogen, cyano, amino, hydroxy, carbamoyl, carboxy,  $C_1-6$  alkanoyl or trifluoromethyl;  
 $n$  is 0 to 4; and

A is  $-(CH_2)_2-$ ,  $-(CH_2)_3-$  or  $-CH=CH-$ ;  
 or a salt thereof.

15 2. A compound according to Claim 1 in which A is  $-(CH_2)_2-$ .

3. A compound according to Claim 1 which is

1-(2-Methylphenyl)-2,3-dihydropyrrolo[3,2-c]cinnoline

1-(2-methylphenyl)-6-methyl-2,3-dihydropyrrolo[3,2-c]cinnoline

1-(2-Methylphenyl)-6-methylpyrrolo[3,2-c]cinnoline

20 1-(4-Hydroxy-2-methylphenyl)-2,3-dihydropyrrolo[3,2-c]cinnoline hydrochloride

N-propyl-2,3-dihydropyrrolo[3,2-c]cinnoline


N-Benzylpyrrolo[3,2-c]cinnoline

1-(4-Hydroxy-2-methylphenyl)-6-methyl-2,3-dihydropyrrolo[3,2-c]cinnoline hydrochloride.

4. A pharmaceutical composition comprising a compound of structure (I) as described in claim 1, or a  
 25 pharmaceutically acceptable salt thereof and a pharmaceutical carrier.

5. The use of a compound of structure (I) or a pharmaceutically acceptable salt thereof in therapy.


6. A process for the preparation of a compound according to Claim 1 which comprises cyclisation of a  
 compound of structure (II)



40 in which  $R^1$  and  $R^2$  are as described for structure (I);  $m$  is 2 or 3 and  $X^1$  and  $X^2$  are leaving groups, in the presence of a compound of structure  $R^3NH_3^+Cl^-$  (III) in which  $R^3$  is as described for structure (I); or

oxidising a compound in which A is  $(CH_2)_2$  to a compound in which A is  $-CH=CH-$ ;  
 forming a salt.

7. A compound of structure (II)



55 in which  $R^1$  and  $R^2$  are as described for structure (I);  $m$  is 2 or 3 and  $X^1$  and  $X^2$  are both halogen, in the presence of a compound of structure  $R^3NH_2$  or  $R^3NH_3^+Cl^-$  (III) in which  $R^3$  is as described for structure (I).



(11) Publication number: **0 380 257 A3**

(12) **EUROPEAN PATENT APPLICATION**

(21) Application number: 90300603.9

(22) Date of filing: 22.01.90

(51) Int. Cl.<sup>5</sup>: **C07D 487/04**, **C07D 471/04**,  
**A61K 31/50**, **C07D 237/28**,  
 //(C07D487/04,237:00,209:00),  
 (C07D471/04,237:00,221:00)

(30) Priority: 23.01.89 GB 8901430

(43) Date of publication of application:  
 01.08.90 Bulletin 90/31

(84) Designated Contracting States:  
 CH DE FR GB IT LI NL

(88) Date of deferred publication of the search report:  
 07.08.91 Bulletin 91/32


(71) Applicant: **SMITHKLINE BECKMAN  
 INTERCREDIT B.V.**  
 28-34 Blaak P.O. Box 2  
 NL-3000 DG Rotterdam(NL)

(72) Inventor: **Brown, Thomas Henry**  
**Smith Kline & French Research Ltd., The  
 Frythe**  
**Welwyn, Hertfordshire AL6 9AR(GB)**  
 Inventor: **Blurton, Peter**  
**Smith Kline & French Research Ltd., The  
 Frythe**  
**Welwyn, Hertfordshire AL6 9AR(GB)**


(74) Representative: **Giddings, Peter John, Dr. et al**  
**SmithKline Beecham, Corporate Patents,**  
**Mundells**  
**Welwyn Garden City, Hertfordshire AL7**  
**1EY(GB)**

(54) Condensed cinnoline derivatives, process and Intermediates for their preparation and  
 pharmaceutical compositions containing them.

(57) Compounds of structure (I)



in which R<sup>1</sup> and R<sup>2</sup> are the same or different and are each hydrogen, C<sub>1</sub>-<sub>6</sub>alkyl, C<sub>1</sub>-<sub>6</sub>alkoxy, phenyl,  
 C<sub>1</sub>-<sub>6</sub>alkylthio, C<sub>1</sub>-<sub>6</sub>alkanoyl, amino, C<sub>1</sub>-<sub>6</sub>alkylamino, diC<sub>1</sub>-<sub>6</sub>alkylamino, halogen or trifluoromethyl;  
 R<sup>3</sup> is hydrogen, C<sub>1</sub>-<sub>6</sub>alkyl or a group



$R^4$  to  $R^6$  are the same or different and are each hydrogen,  $C_1$ - $6$ alkyl,  $C_1$ - $6$ alkoxy,  $C_1$ - $6$ alkylthio, halogen, cyano, amino, hydroxy, carbamoyl, carboxy,  $C_1$ - $6$ alkanoyl or trifluoromethyl;  $n$  is 0 to 4; and A is  $-(CH_2)_2-$ ,  $-(CH_2)_3-$  or  $-CH=CH-$ ; processes for their preparation, pharmaceutical compositions containing them and their use in therapy as inhibitors of gastric acid secretion.



European  
Patent Office

## EUROPEAN SEARCH REPORT

Application Number

EP 90 30 0603

### DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
A	US-A-3 239 526 (SEARLE) * Claim 1; column 1, lines 61-64 * - - -	1,4	C 07 D 487/04 C 07 D 471/04 A 61 K 31/50
P,A	EP-A-0 307 078 (SMITHKLINE) * Claim 1; page 7, lines 48-51 * - - - - -	1,4	C 07 D 237/28 // (C 07 D 487/04 C 07 D 237:00 C 07 D 209:00 ) (C 07 D 471/04 C 07 D 237:00 C 07 D 221:00 )
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			C 07 D 487/00 C 07 D 471/00 A 61 K 31/00 C 07 D 237/00
Place of search		Date of completion of search	Examiner
The Hague		13 May 91	ALFARO I.
CATEGORY OF CITED DOCUMENTS			
X: particularly relevant if taken alone		E: earlier patent document, but published on, or after the filing date	
Y: particularly relevant if combined with another document of the same category		D: document cited in the application	
A: technological background		L: document cited for other reasons	
O: non-written disclosure		&: member of the same patent family, corresponding document	
P: intermediate document			
T: theory or principle underlying the invention			